

# Study of the contribution of the ${}^7\text{Be}(d, p)$ reaction to the ${}^7\text{Li}$ problem in the Big-Bang Nucleosynthesis

Saturday, 20 August 2022 13:45 (15 minutes)

Our research goal is to measure the cross-section of the  ${}^7\text{Be}(d, p)$  reaction in search of a solution to the cosmological  ${}^7\text{Li}$  problem (CLP). The CLP is the overestimation of primordial  ${}^7\text{Li}$  abundance in the standard Big-Bang nucleosynthesis (BBN) model compared to observed abundances, a major unresolved problem in modern astrophysics. A recent theoretical BBN model emphasized the primordial  ${}^7\text{Li}$  abundance is about three times larger than the recent precise observation [1], [2].  ${}^7\text{Li}$  nuclei were considered to be produced predominantly by the electron capture decay of  ${}^7\text{Be}$  after the termination of nucleosynthesis in the standard BBN model. We focus on the  ${}^7\text{Be}(d, p)$  reaction since it is considered one of the contributors to  ${}^7\text{Be}$  destruction in the BBN [3]. We developed a method to produce  ${}^7\text{Be}$  (half life = 53.22 days) target to measure the reaction cross-section in normal kinematics. The experiment was performed at the Tandem Electrostatic Accelerator, Kobe University [4]. A 2.36 MeV proton beam irradiated a natural-Li target to transmute  ${}^7\text{Li}$  particles to  ${}^7\text{Be}$  particles via the  ${}^7\text{Li}(p, n){}^7\text{Be}$  reaction [5]. We produced  $3.03 \times 10^{13}$   ${}^7\text{Be}$  particles in the target after two days of proton irradiation. After the target production, the beam ion was changed to deuterons and the  ${}^7\text{Be}(d, p)$  reaction measured at energies 0.6, 1.0, and 1.6 MeV. The outgoing protons were measured by layered-silicon telescopes placed at 30 and 45 degrees. In this talk, I will report the experimental setup and preliminary results of this study, including the  ${}^7\text{Be}(d, p)$  cross-section.

\\

References

- [1] R. H. Cyburt *et al.*, J. Cosmol. Astropart. Phys. **11**, 012 (2008).
- [2] Brian D. Fields *et al.*, J. Cosmol. Astropart. Phys. **03(2020)010**.
- [3] S. Q. Hou *et al.*, Phys. Rev. C **91**, 055802 (2015).
- [4] "Kobe University Tandem Electrostatic Accelerator" [url{https://www.maritime.kobe-u.ac.jp/en/study/tandem\\_e.html}](https://www.maritime.kobe-u.ac.jp/en/study/tandem_e.html) (Accessed 4th August 2022)
- [5] K. K. Sekharan *et al.*, Nucl. Instr. Meth. **133**, 253-257 (1976).

**Primary author:** INOUE, Azusa (Research Center for Nuclear Physics, Osaka university)

**Co-authors:** TAMII, Atsushi (Research Center for Nuclear Physics, Osaka university, Institute of Radiation Science, Osaka University); CHAN, Phaikying (Research Center for Nuclear Physics, Osaka university); HAYAKAWA, Seiya (Center for Nuclear Study, University of Tokyo); KOBAYASHI, Nobuyuki (Research Center for Nuclear Physics, Osaka university); MAEDA, Yukie (Faculty of Engineering, University of Miyazaki); NONAKA, Kotaro (Faculty of Engineering, University of Miyazaki); SHIMA, Tatsushi (Research Center for Nuclear Physics, Osaka university); SHIMIZU, Hideki (Center for Nuclear Study, University of Tokyo); TRAN, Dinh Trong (Research Center for Nuclear Physics, Osaka university, Vietnam Academy of Science and Technology); WANG, Xuan (Research Center for Nuclear Physics, Osaka university); YAMAGUCHI, Hidetoshi (Center for Nuclear Study, University of Tokyo, National Astronomical Observatory of Japan); YANG, Lei (Center for Nuclear Study, University of Tokyo, China Institute of Atomic Energy); YANG, Zaihong (Research Center for Nuclear Physics, Osaka university, School of Physics, Peking University)

**Presenter:** INOUE, Azusa (Research Center for Nuclear Physics, Osaka university)

**Session Classification:** Young Scientist Session 1