Contribution ID: 23

AA pairing interactions and correlations in multistrange hypernuclei

Thursday, 22 August 2019 15:55 (15 minutes)

We study multistrange Ca, Sn and Pb hypernuclei with $\Lambda\Lambda$ pairing correlations by using the multidimensionallyconstraint relativistic Hartree-Bogoliubov (MDC-RHB) model. The axial deformation is allowed and the $\Lambda\Lambda\omega$ tensor coupling is included to reproduce the small spin-orbit splittings for Λ hyperon. The separable pairing force of finite-range form is used for the pp channel and the ratio of the $\Lambda\Lambda$ pairing to NN pairing strength is determined from the quark model. We find that the shell structure for Λ is very different from that in normal nuclei because of the small spin-orbit splittings in the single Λ spectrum. The pairing energy is similar with HFB calculation results. The $\Lambda\Lambda$ pairing makes the Λ density distribution more symmetric but its influence on the total density distribution can be neglected.

Primary authors: Ms RONG, Yu-Ting (Institute of Theoretical Physics, Chinese Academy of Sciences); Prof. ZHAO, Pengwei (School of Physics, Peking University); Prof. ZHOU, Shan-Gui (Institute of Theoretical Physics, Chinese Academy of Sciences)

Presenter: Ms RONG, Yu-Ting (Institute of Theoretical Physics, Chinese Academy of Sciences)

Session Classification: Young Scientist Session 1